

ORIGINAL PAPERS

Adv Clin Exp Med 2012, 21, 5, 581–589
ISSN 1899–5276

© Copyright by Wrocław Medical University

LUIS A. ROSARIO^{1, A-F}, DANIEL F. ARENCIBIA^{2, A-F}, YOLANDA E. SUAREZ^{3, C, E, F},
JUAN F. INFANTE^{2, C, E, F}, BÁRBARA Y. VALDÉS^{2, B, C}, NIURKA BATISTA^{2, A-F}**Cross-Protection Among Unrelated *Leptospira* Pathogens Serovars: An Unfinished Story****Odporność krzyżowa u niepowiązanych serotypów *Leptospira* – niedokończona opowieść**¹ Institute of Pharmacy and Food Science, University of Havana, Havana, Cuba² Finlay Institute, Havana, Cuba³ Veterinary Medicine Faculty, Agrarian University of Havana (UNAH), Mayabeque, Cuba

A – research concept and design; **B** – collection and/or assembly of data; **C** – data analysis and interpretation; **D** – writing the article; **E** – critical revision of the article; **F** – final approval of article; **G** – other

Abstract

Background. The World Health Organization considers leptospirosis the most neglected zoonotic disease in the last decades. One of the major obstacles in the developing of vaccines for the prevention of leptospirosis is the absence of cross-protection among unrelated serovars. It is accepted that cross-protection among related serovars is due to antibodies generated against lipopolysaccharides (LPs), whereas a cross-protection among unrelated serovars is rarely observed.

Objectives. The objective of the study was to ascertain the existence of cross-protection among vaccine strains of different serovars.

Results. The results of this research demonstrated that a cross-protection among unrelated *Leptospira* serovars strain is possible. The Canicola strain is able to induce protection against homologous, Ballum and Copenhageni strains. The Mozdok strain induced protection only against a homologous challenge. Other strains showed a moderate cross-protection against a heterologous challenge.

Conclusions. These findings suggest that the Canicola and Mozdok strains are ideal candidates for developing a new vaccine formulation for use in Cuba (*Adv Clin Exp Med* 2012, 21, 5, 581–589).

Key words: *Leptospira*, leptospirosis, cross-protection, vaccine, *Mesocricetus auratus*.

Streszczenie

Wprowadzenie. Światowa Organizacja Zdrowia uważa leptospirozę za najbardziej lekceważoną chorobę odzwierzęcą w ostatnich dziesięcioleciach. Jedną z głównych przeszkód w rozwoju szczepionek zapobiegających leptospirozie jest brak odporności krzyżowej wśród niepowiązanych serotypów. Przyjmuje się, że odporność krzyżowa wśród powiązanych serotypów wynika z przeciwciał wytworzonych przeciwko lipopolisacharydom (LPs), a wśród niepowiązanych serotypów odporność krzyżowa występuje rzadko.

Cel pracy. Ustalenie, czy odporność krzyżowa występuje między szczepionkami opartymi na szczepach różnych serotypów.

Wyniki. Badania wykazały, że odporność krzyżowa wśród niepowiązanych szczepów serotypów *Leptospira* jest możliwa. Szczep *L. canicola* jest w stanie wywołać odporność przed szczepami homologicznymi, *Ballum* i *Copenhageni*. Szczep *L. mozdok* wywołał odporność tylko przed homologicznym szczepem. Inne szczepy wykazały umiarkowaną odporność krzyżową przed heterologicznym wyzwaniem.

Wnioski. Wyniki badań sugerują, że szczepy *L. canicola* i *L. mozdok* są idealnymi kandydatami na składnik nowej szczepionki do użytku na Kubie (*Adv Clin Exp Med* 2012, 21, 5, 581–589).

Słowa kluczowe: *Leptospira*, leptospiroza, odporność krzyżowa, szczepionka, *Mesocricetus auratus*.

Leptospirosis is a widespread disease initially described as Weil's syndrome [1]. It is mainly an occupational disease which affects humans and domestic animals, particularly dogs, cattle and swine [2, 3]. Susceptible animals acquire the infection by direct or indirect contact with the urine or tissues of infected animals [3]. *Leptospira* enter the body through small cuts or abrasions, via mucous membranes such as the conjunctiva or through wet skin. They circulate in the blood stream, with the bacteremic phase lasting for up to 7 days [4]. The primary lesion is damage to the endothelium of small blood vessels, leading to localized ischemia in organs, resulting in renal tubular necrosis, hepatocellular and pulmonary damage, meningitis, myositis and placentitis [5].

Leptospira are spirochetes, about 0.1 μm in diameter by 6–20 μm in length [6]. These include both saprophytic and pathogenic species comprising the genus *Leptospira*, which belongs to the family Leptospiraceae, order Spirochaetales. It has over 200 pathogenic serovars, and is divided into 25 serogroups; many different strains with small antigenic differences can be found in some serovars [5, 7]. According to the genetic classification, 13 pathogenic and 6 saprophytic species have been described [5]. The phenotypic and genetic classification do not necessarily coincide. Various molecular methods can be used to differentiate pathogenic and saprophytic leptospira. The most widely used is polymerase chain reaction (PCR) with primers recognizing the gene sequence *lipL32* (the lipoprotein component of the outer membrane), a typical marker present only in pathogenic *Leptospira* [8].

Immunity to leptospirosis is predominantly humoral in humans and most animal species, including dogs, pigs, guinea pigs and hamsters. Antibodies against leptospira lipopolysaccharide (LP) molecules can transfer passive immunity between animals in some models of leptospirosis [9, 10]. Pathogenic *Leptospira* are resistant to the bactericidal activity of normal serum, and in absence of specific antibodies they are neither phagocytized nor destroyed by macrophages [11]. The immune response is also implied in the pathogenesis of leptospirosis through the formation of the immunological complex, the liberation of cytokines and the generation of auto-immune vasculitis [11, 12]. Signs and symptoms of lung, renal and hepatic commitment thus appear in the immune phase when the specific antibodies begin to be detected [12]. Until recently, cross-protection was thought to be restricted to antigen related serovars, since immunity against leptospira LPs following infection is generally highly serovar specific (but not exclusively so) [5, 13]. Some studies carried out in last decade have demonstrated in an indirect

way that cross-protection can exist among serovars of different serogroups [5, 13, 14]. The objective of the current study was to ascertain the existence of cross-protection among vaccine strains of different serovars, all of them of epidemiologic interest for Cuba.

Material and Methods

Bacterial Strains

The *Leptospira spp.* serovars used in this study (Table 1) were cultivated at 28°C in Ellinghausen-McCullough-Johnson-Harris (EMJH) medium supplemented with 10% inactivated rabbit serum. The strains and samples were supplied by the Quality Laboratories of the Finlay Institute, Havana, Cuba. To determine the specificity of the primers, a number of microorganisms were tested, including *L. interrogans* serovar Canicola, *L. biflexa* serovar Patoc (American Type Culture Collection [ATCC] strain) and *Treponema pallidum* spp (ATCC strain).

Table 1. Phenotype classification and LD₅₀ of *Leptospira spp* strains

Tabela 1. Klasyfikacja fenotypowa szczepów LD₅₀ *Leptospira spp.*

Strain (Szczep)	Phenotypic complex/ serovar (Kompleks fenotypowy/ serotyp)	LD50 [UFC/ mL]
FoBa	<i>Leptospira interrogans</i> Ballum	7
CE1	<i>Leptospira interrogans</i> Canicola	4
CE2	<i>Leptospira interrogans</i> Mozdok	6
CE3	<i>Leptospira interrogans</i> Copenhageni	8

The Source and Care of the Hamsters Used in the Study

Golden Syrian hamsters (*Mesocricetus auratus*) were obtained from CENPALAB, Havana, Cuba. They were housed and bred at the Animal House facilities of the Finlay Institute. Ethical approval for the study was obtained from the Ethics Committee of the Finlay Institute.

PCR for Confirming *Leptospira* Pathogenicity

The PCR method as described by Gravekamp et al. [8] for the amplification of *lipL32* was used, with minor variations, to confirm the pathogenicity of the strains used in the study. The vaccine strain *L. interrogans* serovar Canicola Hond Utrecht IV, *L. biflexa* serovar Patoc (ATCC) and *Treponema pallidum* spp (ATCC) were used as controls. For the DNA extraction the cultures were diluted 1-fold in 1 mL of phosphate-buffered saline (PBS: 0.02 mol/L Na₂HPO₄; 0.15 mol/L NaCl, pH 7.2) and centrifuged at 14,000 g for 10 minutes. The pellet was resuspended in 50 µL of PBS and boiled for 10 minutes at 100°C. After processing, the samples were stored at -20°C until they were used for PCR. The *Leptospira interrogans lipL32* sequence was obtained from GenBank, and PCR primers were designed using vector NTI 5.0 (Informax Inc., Bethesda, Maryland, USA). The PCR primers *lipL32* [F] 5'-CGCTGAAATGGGAGTTCGTATGATT-3' and [R] 5'-CCAACAGATGCAACGAAAGATCC TTT-3' were selected, resulting in a 423 bp amplicon. The PCR products (8 µL) were revealed by means of agarose electrophoresis in 2% gel colored with ethidium bromide (0.05 µg/ µL) in buffer TBE 1X.

LD₅₀ Determination

The hamsters (9 weeks old) were divided into 7 groups, each containing 8 hamsters. Six of the groups were given different concentrations of inocula (10⁰, 10¹, 10², 10³, 10⁴ and 10⁵ leptospires) intraperitoneally in 1 ml quantities, as described by Silva et al. [15]. The control animals were administered with the EMJH media used to grow the leptospires, while the inoculated groups were given the leptospires suspended in their respective growth media. The animals were monitored twice daily for up to 21 days post-challenge, but were euthanized whenever signs of terminal disease appeared (severe dehydration, anorexia and immobilization). From the results of the LD₅₀ testing, the concentrations required for the vaccines were obtained using the Reed-Muench method [16].

Vaccine Preparation

To generate the monovalent Ballum vaccines the production methodology developed for vax-SPIRAL® [17] was used with slight modifications. The selected isolates were propagated for three weeks in EMJH medium until dense cultures were observed, using dark-field microscopy (an Olympus BX51 microscope with a dark-field con-

denser). The leptospires were washed three times by centrifugation at 10 000 g for 30 minutes and resuspended in 40 ml of phosphate buffered saline (PBS). After the final wash 0.5% neutral buffered formalin was added for 20 minutes, and the pellet was then similarly washed in PBS four times. The suspension was then diluted to a concentration of 6 × 10⁸ cells/ml. Aluminium hydroxide (1.0 mg/ml) was added as adjuvant and tiomersal (0.05 mg/ml) was added as preservative [18]. The vaccine produced was inoculated onto semi-solid EMJH medium to confirm that there were no viable leptospires, and a loopful was inoculated in a blood agar and broth tioglicolate medium that was incubated at 37°C for 24 hours to confirm the absence of aerobic and anaerobic microorganisms. Monoclonal antibody testing (MAT) was then used to verify that the strains were pure and of their original identities [19]. The vaccine preparations that were found to have satisfactory results in all the tests were packed aseptically in volumes of 5.2 mL, using 10 mL glass bulbs with rubber covers and protective metallic shields and were stored at 4°C until they were used.

The Experimental Challenge and Post-Mortem Examinations

For the vaccine trial, a total of four groups, each composed of 50 hamsters (aged 4 weeks) were used (Table 2). The groups inoculated with a monovalent formulations were each intramuscularly administered a series of two inoculations, six weeks apart. The challenge trial was carried out as described in Table 2. The animals were challenged intraperitoneally with 100 LD₅₀ of each of the isolates.

The animals were monitored twice daily for up to 21 days post-challenge for any sign of illness such as external hemorrhage, dehydration, ruffled hair, decreased activity and isolation from other hamsters in the cages. The clinical signs were scored, using the following scoring system: 0 = no clinical signs; 1 = mild signs such as decreased activity; 2 = moderate signs, including anorexia and depression; 3 = severe clinical signs, including the inability to move or death. Post-mortem examinations were performed and the liver and kidneys were harvested to prove Koch's postulates [20].

Antibody Measurements

Blood was collected by venipuncture of the lateral saphenous vein [21] using microcapillary tubes before each inoculation, prior to the challenge, 2 weeks after inoculation, and then at death or 28 days post-inoculation at euthanasia. Blood collected from hamster was allowed to clot over-

Table 2. The groups of hamsters used in the experiment and the results of tissue-culture and mortality observed in each group**Tabela 2.** Grupy chomików użytych w eksperymencie, wyniki hodowli tkankowej i zaobserwowana śmiertelność w każdej grupie

Group name (Nazwa grupy)	Description (Opis)	Number in group pre-challenge (Liczba osobników w grupie przed badaniem)	Mortality (Śmiertelność) %	# culture positive/ # livers evaluated (Pozytywna hodowla/ liczba ocenionych preparatów wątroby)	# culture positive/ # kidneys evaluated (Pozytywna hodowla/ liczba ocenionych preparatów nerek)	Category of pathological findings (Rodzaj patologicznych zmian)
A	immunized with Canicola	40				
	challenge with Canicola	10 each	0	0/10	0/10	0
	challenge with Mozdok		100	10/10	10/10	2
	challenge with Copenhageni		0	0/10	0/10	0
	challenge with Ballum		0	0/10	0/10	0
B	immunized with Mozdok	40				
	challenge with Canicola	10 each	100	10/10	10/10	2
	challenge with Mozdok		0	0/10	0/10	0
	challenge with Copenhageni		100	10/10	10/10	2
	challenge with Ballum		100	10/10	10/10	1
C	immunized with Copenhageni	40				
	challenge with Canicola	10 each	60	3/10	5/10	1
	challenge with Mozdok		100	10/10	10/10	2
	challenge with Copenhageni		0	0/10	0/10	0
	challenge with Ballum		20	2/10	2/10	1
D	immunized with Ballum	40				
	challenge with Canicola	10 each	0	2/10	3/10	1
	challenge with Mozdok		100	8/10	10/10	2
	challenge with Copenhageni		30	2/10	3/10	1
	challenge with Ballum		0	0/10	0/10	0

Category 0: no pathology. Category 1: subcutaneous jaundice, bleeding of the nail beds, hemorrhages of the kidney, liver is friable with discolorations and smooth edges. Histologically there is moderate nephritis, hepatocyte necrosis and cellular dis cohesion. Category 2: gross pitting on the surface of the kidney, with capsular adhesions. Histologically moderate nephritis with acute renal damage to epithelial cells and tubular necrosis. With respect to the liver, there is increased binucleation of hepatocytes, and an increase in the number of cells with foamy cytoplasm.

Kategoria 0: bez patologii. Kategoria 1: żółtaczka podskórna, krwawienie z macierzy pazura, krwotoki z nerki, krucha wątroba z przebarwieniami i gładkimi zakończeniami. Histologicznie umiarkowane zapalenie nerek, martwica hepatocytów i dekohezja komórek. Kategoria 2: wyrwy na powierzchni nerki, zrosty otoczkowe. Histologicznie umiarkowane zapalenie nerek z ostrą uszkodzeniem nabłonka i martwicą kanalików. W wątrobie, istnieje zwiększona liczba jąder w hepatocytach i zwiększenie liczby komórek z „piankową” cytoplazmą.

night and centrifuged at 700 g for 20 minutes; the serum removed and stored at -20°C until analysis. For the ELISA test, 96-well ELISA plates (MaxiSorp, Nunc, Denmark) were coated using 50 μl per well of 5 $\mu\text{g/ml}$ of each serovar sonicate diluted

in 0.1M Na_2HPO_4 (pH 9.0). The plates were incubated overnight at 4°C , washed with PBS containing 0.05% Tween 20 (PBST), blocked at room temperature for 3 hours with 200 μl of 1% BSA in PBS, and again washed with PBST. Serum samples

were thawed and diluted 1:100 and 1:1000 with 1% BSA in PBST, then added to the plates at 100 μ l per well and incubated at room temperature for 2 hours. These concentrations were found to give readings in the linear part of the assay without increasing background readings in pilot experiments that evaluated a large range of serum dilutions from 1:10 to 1:10⁶ and included known negative and known positive sera. After washing, 100 μ l of mouse anti-bovine IgG conjugated to horseradish peroxidase (1 μ g/ml, Serotec) diluted with 1% BSA in PBST was added to the plates and incubated at room temperature for 45 minutes. Following 10 μ l of 30% H₂O₂ was added to 11 ml of freshly thawed 0.3 mg/ml 2,2'-Azino-bis-(3-ethylbenzthiazoline-6-sulfonic acid) (ABTS) in 0.1M citric acid (pH 4.35) and 100 μ l per well was added to the plates. The plates were incubated with the substrate at room temperature for 25 minutes, the color reaction was stopped by adding 50 μ l per well of 1% SDS, and optical density at 405 nm was determined using a Dynex plate reader. The microscopic agglutination test was conducted as described by Cole et al. [22] and the sera were serially diluted until an endpoint titer was obtained, as is conventional. Antibody levels were therefore expressed as IgG U/ mL.

Statistical Issues and Analyses

The data was analyzed at a 5% significance level using STATISTICA software version 6.1 (USA). Kaplan-Mayer tests were performed to determine if there were significant differences in survival rates.

Results

PCR *Leptospira* Pathogenicity

The four strains of *Leptospira* spp. used for the study tested positive by PCR. Figure 1 shows an ethidium bromide stained agarose gel after electrophoresis in which a 423 base pair amplicon represents the *lipL32* sequence, which is a feature of pathogenic *Leptospira* genus. Besides the control strain *Leptospira interrogans* Canicola Hond Utrecht IV produced a 423 base pair amplicon, which was absent from PCR products obtained from *L. biflexa* serovar Patoc (ATCC) and *Treponema pallidum* spp (ATCC).

LD₅₀ Determination

All the *lipL32* positive isolates showed a high virulence. These strains caused death within 6–14 days of inoculation. The Canicola strain tested



Fig. 1. *lipL32* polymerase chain reaction. Line 1: Weight marker [4521 Promega], Lines 2–3: *L. interrogans* Canicola serovar, Lines 4–5: *L. interrogans* Mozdok serovar, Lines 6–7: *L. interrogans* Copenhageni serovar, Lines 8–9: *L. interrogans* Ballum serovar, Line 10: *L. interrogans* Canicola (ATCC) serovar, Lines 11–12: *L. biflexa* Patoc (ATCC) serovar, Line 13: *Treponema pallidum* spp (ATCC), Line 14: Weight marker 720–20 pb.

Ryc. 1. *lipL32* reakcja łańcuchowa polimerazy. Linia 1: oznaczenie masy [4521 Promega]. Linia 2–3: *L. interrogans* serotyp canicola. Linia 4–5: *L. interrogans* serotyp Mozdok. Linie 6–7: *L. interrogans* serotyp Copenhageni. Linie 8–9: *L. interrogans* serotyp Ballum. Linia 10: *L. interrogans* serotyp canicola (ATCC). Linia 11–12: *L. biflexa* serotyp Patoc (ATCC). Linia 13: *Treponema pallidum* spp. (ATCC). Linia 14: oznaczenie masy 720–20 pb

caused death within 8–10 days of inoculation. Mozdok and Copenhageni caused death within 7–10 days, while Ballum caused death within 6–16 days. The LD₅₀ value of Canicola, Mozdok, Copenhageni and Ballum were 11, 9, 8 and 10 organisms respectively. The statistical comparison of these results did not show a significant difference among them.

Antibody Measurements

The IgG response generated by the immunization (two doses) with the different monovalent formulations against the Canicola, Mozdok, Copenhageni and Ballum serovars are shown in Figure 2. In all the groups the antibodies' response against the heterologous serovars were similar to the negative controls and pre-immune values. The IgG response in the group immunized with the Canicola monovalent was revealed in 70% of the animals. After applying the second dose, the values of IgG generated were very significant relative to the values obtained after the first dose. Also, after the first dose 100% of the animals in the immunized groups were positive. A similar response was observed with other monovalents; the main differences were in the percents of responder animals after the first dose (60–80%) and intensity of response. The more immunogenic strains were Canicola and Ballum, and in all groups the IgG response were detected until death or euthanized.




Fig. 2. IgG response induced in hamsters against *Leptospira* serovars after the immunization with two doses of monovalent formulations [A] Canicola, [B] Mozdok, [C] Copenhageni, [D] Ballum

Ryc. 2. Odpowiedź IgG wywołana u chomików przeciwko serotypom *Leptospira* po szczepieniu dwoma dawkami formacją jednowartościową a) *canicola*, b) *Mozdok*, c) *Copenhageni*, d) *Ballum*

Experimental Challenge and Post-Mortem Examinations

Figure 3 summarizes the survival rate of vaccinated and unvaccinated animals challenged with the different strains. Hamsters vaccinated with the experimental monovalent formulations and later challenged with a homologous strain showed no clinical signs of leptospirosis and there was no post-challenge mortality. These animals were also culture-negative, with no gross or histological pathological lesion. The same results were obtained in animals vaccinated and not challenged, used as safety controls for each of the monovalent formulations. The unvaccinated challenged hamsters displayed the greatest morbidity (an average histopathological score of 2) with 100% of the hamsters showing clinical signs and 100% mortality.

Mortalities in vaccinated animals challenged against heterologous strains occurred after the vaccination period. For the groups of hamsters inoculated with a Ballum monovalent formulation and challenged with Mozdok and Copenhageni, severe clinical signs were observed in 60–100% of the animals between 8 and 20 days post-challenge, giving mortality rates of 100% and 70% respectively; in

contrast, the animals challenged with Canicola did not show mortality or clinical signs. The animal groups inoculated with Canicola monovalent formulation and challenged with Copenhageni and Ballum showed a 100% survival rate without clinical signs, but the animals in this group challenged with Mozdok did not survive: They showed 100% mortality, occurring between 6 and 18 days post-challenge, with very mild clinical signs (average histopathological score = 1). The Copenhageni formulation partially protects against a Canicola challenge (80%), while it protects against Ballum and Mozdok only 30% and 0% respectively (average histopathological score = 2). Finally, the animals immunized with a Mozdok monovalent formulation and challenged with heterologous strains showed 100% mortality; death occurred between 10 and 23 days (average histopathological score = 2).

The organ culture results and gross histopathological lesions from the animals that died or were euthanized are summarized in Table 2. The most severe lesions included hemorrhages of the kidneys and liver. The major histopathological lesions at the kidney level displayed renal tubular necrosis with occasional polymorphonuclear cell infiltration and loss of normal liver architecture.




Fig. 3. Percentage of surviving hamsters immunized with monovalent formulations [A] Canicola, [B] Mozdok, [C] Copenhageni, [D] Ballum 21 days post-challenge with 100 DL_{50} homologous and heterologous strains

Ryc. 3. Odsetek przeżywających chomików immunizowanych jednowartościowym preparatem a) *canicola*, b) *Mozdok*, c) *Copenhageni*, d) *Ballum* 21 dni po prowokacji 100 szczepami DL_{50} homologicznie i heterologicznie

Discussion

To test the hypothesis that cross-protection can be established among unrelated serovars of pathogenic *Leptospira* strains, the authors evaluated the capacity of whole cellular monovalent formulations to confer protection against a lethal challenge in hamsters. The selection of the strains for this assay was based on their epidemic importance for Cuba. All the isolates tested were positive for the *lipL32* gene, which is found in pathogenic *Leptospira*, and also showed high levels of virulence in Golden Syrian hamsters, which are the animal model most frequently [23] because acute infections exhibit a similar symptomatology to the human infectious process [24, 25]. The virulence of the strains used is an important characteristic to assure the accuracy of the results, because several differences have been observed in leptospirosis pathogenesis using virulent and nonvirulent isogenic strains [26–29]. Extracellular proteins, external membrane lipoproteins and adhesins are expressed more by virulent strains than nonvirulent ones [26, 27]. This unequal antigen architecture translates into a different protector capacity [27]. The LD_{50} experiment results showed that hamster death occurred between 7 and 16 days

post-challenge. The timing of the deaths and clinical signs observed were consistent with those of other LD_{50} experiments in hamsters using these serovars [14].

The homologous and heterologous challenge of immunized hamsters with monovalent formulations of inactivated whole cells with these virulent strains evidenced the existence of a complex immune response able to establish cross-protection only among some of the studied serovars, establishing in most of the cases only a one-way protection – that is, the protection conferred by one serovar against another is not necessarily a reciprocal relation. Immunization with the Canicola serovar establishes protection against the rest of the serovars, with exception of Mozdok; on the other hand, immunization with Copenhageni does not guarantee complete protection against a challenge with Canicola. Similarly, immunization with Ballum protects the animals against a challenge with Copenhageni in 80% of the cases, while immunization with Copenhageni only protects the animals challenged with Ballum in 40% of the cases.

Another element of interest in this study is the absence, in all cases, of IgG response against the heterologous serovars, including those in which a partial or total cross-protection was observed.

Some of the causes that can explain high protection levels in the absence of specific IgG antibodies may be the presence of protective IgM antibodies and/or the influence of the cellular response [5]. In Golden Syrian hamsters the presence of IgM antibodies is more relevant, because in this species the immune response against leptospirosis is predominantly humoral [5] and it is characterized as directed fundamentally against LPs [28]. The protection conferred by LPs is serovar specific [29]; in contrast, the response against external membrane protein is related to cross-protection among unrelated serovars [30]. During a natural infection the humoral response against membrane proteins is of much lower relevance than LPs, possibly due

to the fact that membrane proteins are present in smaller quantities than LPs and are less accessible to the immune system. The major consequence of this is a lack of long-term protection (IgG antibody response) and a marked decrease in the cross-protection [5, 29, 30].

The obtained results confirm that the cross-protection in *Leptospira* is a complex phenomenon that depends, among other factors, on 1) the serological characteristic of the evaluated strain, and 2) the host immunology system. The authors suggest further evaluation of the Canicola and Mozdok strains as candidates for developing a new whole vaccine formulation capable of generating a broad heterologous protective spectrum.

Acknowledgements

The authors wish to thank the Research Vice-Presidency of Finlay Institute for funding the project, as well as the technicians at the Animal House at Finlay Institute and G. Sierra Ph.D. for their assistance.

References

- [1] **Vinetz JM:** Leptospirosis. *Curr Opin Infect Dis* 2001, 14, 527–538.
- [2] **Trueba G, Zapata S, Madrid K, Cullen P, Haake D:** Cell aggregation: a mechanism of pathogenic *Leptospira* to survive in fresh water. *Int Microbiol* 2004, 7, 35–40.
- [3] **Waitkins SA:** Leptospirosis as an occupational disease. *Br J Ind Med* 1986, 43, 721–725.
- [4] **McBride AJ, Athanazio DA, Reis MG, Ko AI:** Leptospirosis. *Curr Opin Infect Dis* 2005, 18, 376–386.
- [5] **Adler B, de la Peña M:** *Leptospira* and Leptospirosis. *Vet Microbiol* 2009, 2, 4382–4392.
- [6] **Faine S, Adler B, Bolin C, Perolat P:** *Leptospira* and leptospirosis. *Medisci Melbourne* 1999, 135–165.
- [7] **Kuriakose M, Eapen CK, Paul R:** Leptospirosis in Kolenchery, Kerala, India: epidemiology, prevalent local serogroups and serovars and a new serovar. *Eur J Epidemiol* 1997, 13, 691–697.
- [8] **Gravekamp C, Van De Kemp H, Franzen M, Carrington D, Schoone GJ, Van Eys GJJM:** Detection of seven species of pathogenic leptospires by PCR using two sets of primers. *J Gen Microbiol* 1993, 139, 1691–1700.
- [9] **Faine S:** *Leptospira* and leptospirosis, CRC Press, Boca Raton [FL], 1994.
- [10] **Jost BH, Adler B, Vinh T, Faine S:** A monoclonal antibody reacting with a determinant on leptospiral lipopolysaccharide protects guinea pigs against leptospirosis. *J Med Microbiol* 1986, 22, 269–275.
- [11] **Levett PN:** Leptospirosis. *Clin Microbiol Rev* 2001, 14, 296–326.
- [12] **Cinco M, Domenis R, Perticarari S, Presani G, Marangoni A, Blasi E:** Interaction of leptospires with murine microglial cells. *New Microbiol* 2006, 29[3], 193–199.
- [13] **Sonrier C, Branger C, Michel V, Ruvoen-Clouet N, Ganiere JP, Andre-Fontaine G:** Evidence of cross-protection within *Leptospira interrogans* in an experimental model. *Vaccine* 2001, 19[1], 86–94.
- [14] **Silva EF, Medieros MA, Mc Bride AJA, Matasunga J, Esteves GS, Ramos JGR:** The terminal portion of leptospiral immunoglobulin-like protein Lig A confers protective immunity against lethal infection in the hamster model of leptospirosis. *Vaccine* 2007, 25[33], 6277–6286.
- [15] **Silva E, Santo C, Athanazio D, Seyffert N, Seixas F, Cerqueira GM:** Characterization of virulent *Leptospira* isolates in hamster model. *Vaccine* 2008, 26, 3892–3896.
- [16] **Reed LJ, Muench H:** A simple method of estimation fifty percent end-points. *Am J Hyg* 1938, 27, [3], 493–497.
- [17] **Finlay Institute:** Sanitary Registration of vax-SPIRAL (trivalent antileptospirosic vaccine), 1998, 23–25.
- [18] **González A, Rodríguez Y, Batista N, Valdés Y, Núñez JF, González M:** Booster effect of an active immunization with *Leptospira interrogans* serogroup Ballum in hamsters vaccinated with vax-SPIRAL. *VacciMonitor* 2003, 12[2], 1–6.
- [19] **KIT Biomedical Research.** International Course on Laboratory Methods for the Diagnosis of Leptospirosis. Amsterdam: Royal Tropical Institute; 2006.
- [20] **Falkow S:** Molecular Koch's postulates applied to microbial pathogenicity. *Rev Infect Dis* 1988, 10[2], 274–276.
- [21] **Hem A, Smith AJ, Solberg P:** Saphenous vein puncture for blood sampling of the mouse, rat, hamster, gerbil, guinea pig, ferret, and mink. *Lab Animal* 1998; 32, [4], 364–368.
- [22] **Cole JR, Sulzer CR, Pursell AR:** Improved microtechnique for the leptospiral microscopic agglutination test. *Appl Microbiol* 1973, 25, 979–980.
- [23] **Oliva R, Infante J, González M, González I, Fariñas M:** Clinical and pathological characterization of leptospirosis in Golden Syrian hamster and Duncan Hartley guinea pig by means of experimental infection with three *L. interrogans* serovars. *VacciMonitor* 1998, 7[5], 8–13.

- [24] **Matsunaga J, Young T, Barnett J, Barnett D, Bolin C, Haake D:** Novel 45-kilodalton leptospiral protein that is processed to a 31-kilodalton growth-phase-regulated peripheral membrane protein. *Infect Immun* 2002, 70[1], 323–334.
- [25] **Barnett J, Barnett D, Bolin C, Summers T, Wagar E, Cheville N:** Expression and distribution of leptospiral outer membrane components during renal infection of hamsters. *Infect Immun* 1999, 67, 853–861.
- [26] **Merien F, Truccolo J, Baranton G, Perolat P:** Identification of a 36-kDa fibronectin-binding protein expressed by a virulent variant of *Leptospira interrogans* serovar Icterohaemorrhagiae. *FEMS Microbiol Lett* 2000, 185, 17–22.
- [27] **Merien F, Baranton G, Perolat P:** Invasion of Vero cells and induction of apoptosis in macrophages by pathogenic *Leptospira interrogans* are correlated with virulence. *Infect Immun* 1997, 65[2], 729–738.
- [28] **Suepaul SM, Carringtonb CV, Campbell M, Bordea G, Adesiyuna AA:** Study on the efficacy of *Leptospira* vaccines developed from serovars isolated from Trinidad and comparison with commercial vaccines using a hamster model. *Vaccine* 2010, 28 [1], 5421–5426.
- [29] **Chapman A, Everard COR, Faine S, Adler B:** Antigens recognized by the human immune response to severe leptospirosis in Barbados. *Epidemiol Infect* 1991, 107, 143–155.
- [30] **Biswas D, Roy S, Vijayachari P, Sugunan A, Nararajaseenivasan K, Sehgal S:** Comparison of immunoreactive proteins of commonly circulating serogroups of *Leptospira* in Andaman Islands. *India J Med Res* 2005, 121, 151–158.

Address for correspondence:

Luis A. Rosario
Institute of Pharmacy and Food Science
University of Havana
222 Street between 25 and 27 Avenues
Lisa Municipality
Havana
Cuba
E-mail: darencibia@cecmecmed.sld.cu
Tel.: 0537 271 11 58

Conflict of interest: None declared

Received: 20.03.2012

Revised: 27.04.2012

Accepted: 8.10.2012